

<i>Product Licence Number</i>	<i>Company Name</i>	<i>Product Name</i>	<i>Active Ingredients</i>	<i>Indications</i>	<i>Date of Authorisation</i>
00041/0049	Bioglan Laboratories Ltd.	Epaderm	Emulsifying Wax 30-000 ww Paraffin Soft Yellow 30-000 ww	For general use as an emollient to moisturise and soften dry skin. Epaderm may be used instead of soap. General Sales List	28th February 1996
00053/0254	Schering Health Care Ltd.	Magnevist	Gadopentetic Acid Dimeglumine Salt 469-010 mg	As a paramagnetic contrast medium in cranial, spinal and whole body magnetic resonance imaging (MRI) and for the evaluation of renal function. Prescription Only Medicine	26th February 1996
00053/0258	Schering Health Care Ltd.	Water for Injection	none	Water for injection is used to dissolve the accompanying Levovist granulate and create a suspension. Prescription Only Medicine	23rd May 1996
00057/0404	Pfizer Limited	Vibramax 100 (Doxycycline Capsules 100 mg)	Doxycycline Hyclate 115-400 mg	(See PL00057/0405) Prescription Only Medicine	1st March 1996
00057/0405	Pfizer Limited	Vibramax 50 (Doxycycline Capsules 50 mg)	Doxycycline Hyclate 57-700 mg	Pneumonia: respiratory tract infections; pneumonia and other lower respiratory tract infections due to susceptible strains of <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> , <i>Klebsiella pneumoniae</i> and other organisms. <i>Mycoplasma pneumoniae</i> pneumonia. Treatment of chronic bronchitis, sinusitis. Genito-urinary tract infections: cystitis, urethritis, pyelonephritis, caused by susceptible strains of <i>Escherichia coli</i> , <i>Neisseria gonorrhoeae</i> , the <i>Klebsiella-Enterobacter</i> group, <i>Staphylococcus</i> species and <i>Chlamydia trachomatis</i> . A single large dose of doxycycline has effectively treated acute gonococcal anterior urethritis in the adult male. However, a more extended therapy elicited the highest cure rates. Soft tissue infections: infected traumatic and postoperative wounds, furunculosis, impetigo, cellulitis, abscess and paronychia, caused by susceptible strains of the <i>Klebsiella-Enterobacter</i> group, as well as susceptible strains of <i>Staphylococcus aureus</i> , <i>Streptococcus</i> species, <i>Escherichia coli</i> and <i>Staphylococcus albus</i> . Doxycycline treatment of soft tissue infections should always be carried out in conjunction with any necessary indicated surgical procedures. Dermatological infections: <i>Acne vulgaris</i> and <i>Acne conglobata</i> . As doxycycline is one of the tetracycline group of antibiotics, it may be expected to be useful in the treatment of infections which respond to other tetracyclines such as: Gastro-intestinal infections: caused by susceptible strains of organisms such as <i>Shigella</i> species, <i>Salmonella</i> species, <i>Entamoeba histolytica</i> and enteropathogenic <i>Escherichia coli</i> . Ophthalmic infections: caused by susceptible strains of <i>Staphylococci</i> , <i>Gonococci</i> and <i>Haemophilus influenzae</i> . Doxycycline is indicated for the treatment of inclusion conjunctivitis either alone or in combination with topical agents. Doxycycline is also suitable in the treatment of trachoma. However, the infectious agent is not eliminated on all occasions, as tested by immunofluorescence techniques. Miscellaneous: trigonitis and prostatitis caused by <i>Proteus</i> species. Psittacosis, certain other infections caused by susceptible strains of <i>Bacteroides</i> species, <i>Listeria</i> species, <i>Rickettsia</i> species, <i>Yersinia</i> species, <i>Brucella</i> species (in combination with streptomycin) and <i>Bordetella pertussis</i> , <i>Calymmatobacterium granulomatis</i> , <i>Spirochaetes</i> ( <i>Treponema</i> species), <i>Bacillus anthracis</i> , <i>Neisseria meningitidis</i> and <i>Clostridium welchii</i> . Doxycycline may be a useful adjunct to amoebicides in the treatment of acute intestinal amoebiasis. Prescription Only Medicine	1st March 1996